

A review of the anomalous origin of the left coronary artery from the anterior sinus of valsalva: Is prevention possible ?

Craig T Nakamura MD*

Danelo R Canete MD**

Jeffrey M Lau MD⁺

The origin of the left coronary artery from the anterior sinus of Valsalva is a rare coronary anomaly. Nevertheless, it remains an important pathological entity because of the possibility of its clinical and surgical consequences. Physicians should therefore be aware of this condition and consider it in their differential diagnosis of ischemic heart disease. This is a case report and review of the literature.

Introduction

The purpose of this paper is to provide an overview and to familiarize physicians with this anomaly. A review of 65 autopsies and cases of sudden death among these patients is included. In addition, we report a case of anomalous origin of the left coronary artery from the anterior sinus of Valsalva in Honolulu, Hawaii.

Case Report

An unmarried, 38-year-old Filipino woman was admitted to the hospital with a long-standing history of angina at rest and upon exercise. She had a history of diabetes, hypertension, and anemia; her only current medication was Nitrostat. Her past medical history included a tubal ligation in 1975 and two bladder infections, the last one in March 1990. Her family history was significant for sudden death—her mother died at age 45 of heart disease and the patient's child died at the age of 1 of an unknown cause.

Her blood pressure was 110/70, pulse 60 and regular, respirations 20, and temperature 98.7°C. The physical examination was otherwise unremarkable. Laboratory findings included a CBC, serum chemistries, serum electrolytes, and urinalysis were within normal limits. Echocardiography revealed mild cardiomegaly with left ventricular enlargement. The patient also had a positive thallium stress-test.

* Second-year medical student
John A Burns School of Medicine
University of Hawaii

** Cardiologist
St. Francis Medical Center

+ Clinical Assistant Professor of Surgery
John A Burns School of Medicine
University of Hawaii

Request for reprints:
Jeffrey M Lau, MD
1329 Lusitana Street, Suite 108
Honolulu, HI 96813
Submitted June 1992

Angiography was performed to rule out coronary artery disease or an equivalent. It revealed normal left ventricular function and a large right coronary artery. There was a small left coronary artery arising from the right sinus of Valsalva that passed between the great vessels and provided primarily a circumflex distribution. (Figure 1: x-ray).

Surgical correction consisted of a coronary artery bypass graft. The left internal mammary artery was anastomosed to the left anterior descending coronary artery and a saphenous vein segment was used to connect with the circumflex system. The patient's postoperative course was uneventful with the exception of a single episode of low serum glucose (46 mg/dl). She was discharged 6 days later on Ecotrin and Persantine.



Figure 1: x-ray

Discussion

Normal coronary arteries have an architecture that is "...observed in at least 1% of unselected cases"¹. This definition allows for *normal variants*; therefore, any variation that occurs in less than 1% is an anomaly¹. The incidence of coronary anomalies has ranged from 0.28% to 1.2%²⁻¹⁰. Coronary anomalies can be divided into 3 categories:

- 1) Associated: A variation in response to a primary cardiac pathology.
- 2) Major: An abnormal coronary artery connection with a cardiac chamber or an abnormal origin from the pulmonary artery.
- 3) Minor: An abnormal origin from the aorta but with a normal distal circulation.

Anomalous origin of the left coronary artery from the anterior sinus of Valsalva fits into the third category. Its inci-

dence has been reported between 0.02% and 0.19%^{2,3,4,9,11,12}. It is considered to be a minor anomaly because the blood entering both coronary arteries is fully oxygenated; this variation is compatible with life. However, numerous studies have documented that it is not such a benign condition after all. Cheitlin reported a 27% risk of sudden death in patients with this anomaly¹³.

The various routes an anomalous left coronary artery can take will be discussed prior to addressing possible mechanisms of these sudden deaths.

There are 4 pathways an anomalous left coronary artery can take as it travels from its origin in the sinus of Valsalva to its final destination by dividing into the left anterior descending and circumflex arteries. The first is known as the retroaortic pathway (Figure 3). The left coronary artery leaves the anterior sinus of Valsalva and courses posteriorly around the aorta to its normal distribution. Presently, no cases of sudden death in patients with this pathway have been reported. In addition, Murphy reported a myocardial infarct over the distribution of the left coronary artery in one such patient. Symptoms were alleviated with bypass surgery of the anomalous left coronary artery although both coronary arteries were devoid of atherosclerosis¹⁴. This strongly suggests that the ischemia was caused by the retroaortic anomaly.

The second course an anomalous left coronary artery can take is called the anteropulmonic pathway (Figure 4). The left

coronary artery leaves the sinus of Valsalva and travels anteriorly to the pulmonary artery to its normal distribution. Roberts reported sudden death in one such patient¹⁵. Pachinger presented a patient with such an anomalous left coronary artery who had angina despite patent coronary arteries¹⁶. Chaitman reported myocardial infarction in the left coronary artery's distribution in another patient with patent coronary arteries². These studies of ischemia in the presence of patent but anomalous coronary arteries point to the possible cause of sudden death. Roberts and Kragel also reported sudden death in one such patient¹⁵. This evidence indicates that the anteropulmonic route of an anomalous left coronary artery is not a benign condition.

The third course is known as the interarterial pathway, in which the left coronary artery courses between the aorta and pulmonary artery (Figure 5). This is the most hazardous form and is responsible for a vast majority of the sudden deaths attributed to an anomalous left coronary artery from the anterior sinus of Valsalva.

There is a fourth, benign type of anomalous left coronary artery that has a septal pathway (Figures 6 and 7). The left coronary artery leaves the aorta and burrows into the ventricular septum, and it emerges anteriorly to branch into the left anterior descending and circumflex arteries which follow their usual courses. In a review of the literature, there has been only one

(Continued) ➤

Legend : All superior views

- CX - Circumflex
- LAD - Left Anterior Descending Artery
- LMC - Left Main Coronary Artery
- RCA - Right Coronary Artery
- PA - Pulmonary Artery
- LSV - Left Sinus of Valsalva
- RSV - Right Sinus of Valsalva




Figure 2: Normal anatomy of the coronary arteries.




Figure 3: Anomalous left main coronary artery coursing posterior to the aorta.




Figure 4: Anomalous left main coronary artery coursing anterior to the pulmonary artery.




Figure 5: Anomalous left main coronary artery coursing between the aorta and pulmonary artery.

A REVIEW OF THE ANOMOLOUS ORIGIN OF THE LEFT CORONARY ARTERY FROM THE ANTERIOR SINUS OF VALSALVA: IS PREVENTION POSSIBLE

(Continued from page 295)




Figure 6: Transverse section through heart showing anomalous LMCA coursing in the ventricular septum view from below..

Figure 6 legend

A	—	Aorta
PT	—	Pulmonary
SVC	—	Superior Vena Cava
IVC	—	Inferior Vena Cava
LMC	—	Left Main Coronary Artery
CX	—	Circumflex Artery
LAD	—	Left Anterior Descending Artery
RVOT	—	Right Ventricular Outflow Tract
RA	—	Right Atrium
RV	—	Right Ventricle
LV	—	Left Ventricle




Figure 7: Anomalous left main artery coursing in the ventricular septum (anterior view).

patient documented with this septal pathway which might have contributed to his death¹⁷. This condition is considered to be relatively benign, and patients with the septal route typically live to advanced ages.

There has been much speculation as to the exact cause of sudden death in patients who have anomalous left coronary arteries originating from the anterior sinus of Valsalva. Most cases have been directly related to exercise. Of 39 cases of sudden death in patients under the age of 30 reviewed in the literature, 34 were exercise-related.

The most popular theory is that the anomalous coronary artery is compressed between the aorta and pulmonary artery during exercise^{3,18,19,30}. Ischemia could be caused by the unsatisfied need for increased oxygen consumption by the myocardium and in peripheral vasodilation, or by an increase in systolic pressure leading to compression of the anomalous coronary artery.

Murphy¹⁴ documented a case in which the density of contrast medium in the anomalous left coronary artery clearly diminished during systole.

This explains why sudden death almost never happens in patients with an anomalous septal left coronary; the buried left artery might be protected from compression.

Cheitlin disagreed with the compression theory and said that the pulmonary artery, a low-pressure system, would be an unlikely candidate in the compression of the left coronary artery¹³. However, patients with normal right ventricles and pulmonary vasculature have been found to develop marked transient pulmonary hypertension both before and after a coronary bypass operation¹¹. Corday and associates have found that pulmonary hypertension can compress the coronary arteries²⁰. Thus, transient pulmonary hypertension with a physiologic increase in systolic blood pressure during exercise could compress the anomalous left coronary and hinder its flow.

Several authors have discovered atherosclerotic stenosis in the portion of the anomalous left coronary artery that passes between the aorta and pulmonary artery. These plaques were postulated to be the result of chronic compression^{7,19}. This is unlikely to be the sole cause of sudden death, but in conjunction with left coronary artery compression could compromise coronary blood flow.

Another theory based on ischemia deals with the acute angle

SURGICAL APPROACH




Figure 8a: Anomalous LMC




Figure 8b: Exposure of the anomalous LMC




Figure 8c: The ostium is split and the incision is extended above the intercoronary commissure until the midpoint of the left coronary.



Figure 8d: The intima of the LMCA and aorta are joined, and the inter-coronary commissure is reattached

present in many cases as the artery leaves the aorta. An acute angle can be defined as an angle of less than 45 degrees between the aorta and the proximal portion of the left coronary artery²¹. Several authors have noted that an acute angle could lead to luminal narrowing and reduced myocardial blood supply from luminal kinking and intramural stretching as cardiac output and pressures are elevated during exercise^{3,7,13,19,22}. Virmani, in a study of sudden death victims, found a significant increase in the incidence of such acute angles when compared with a control group²¹; acute angles were present in 18 of the cases reviewed²¹. A slit-like ostium was commonly found in association with an acute angle of the anomalous left coronary artery, often in conjunction with a flap-like closure. This condition was found in 11 of the cases reviewed, 10 of which possessed an acute angle. The already narrowed ostium could be narrowed further with exercise as the aorta distends, reducing coronary blood supply.

In addition, several authors have reported that the proximal left coronary artery that exits the ostium travels within the wall of the aorta, sharing a common intima^{23,24}. With exercise, the aorta dilates, and as Sacks proposed, the intramural portion becomes flattened, reducing blood flow²³.

Virmani also described ostial ridges (defined as a ridge whose surface area occupied 50% of the coronary ostial area). In his study, 22 sudden death victims were compared with a control group of 19 patients who died of known causes. All 41 hearts were examined for abnormalities such as acute angle takeoff and ostial valve-like ridges. Patients with the anomalous left coronary artery with origin from the anterior sinus of Valsalva had a significantly higher incidence of ostial ridges than their control group counterparts. The ridges were thought to impede flow by compressing the left coronary artery as the aorta dilated during exercise²¹.

Another possible cause for ischemia is congenitally hypoplastic coronary arteries. This is not the sole factor leading to sudden death in patients with anomalous left coronary arteries because only 5 of the cases possessed a left coronary artery significantly smaller than the

(Continued) ►

Where can **YOU GO** this weekend?

... **to explore a rain forest,**
... **ocean kayak,**
... **visit ancient temples,**
... **shoot a round of golf**
... and still be back
in your office on
Monday morning?

MOLOKA'I

You make it a Moloka'i weekend.
Kaluakoi makes it affordable. We have
packages from as little as \$99*.
And you'll have the time of your life.

For Reservations/Information
call your travel agent or call
Kaluakoi Hotel

TOLL FREE 1-800-435-7208

Kaluakoi Hotel & Golf Club • Moloka'i, HI 96770 • Managed by Colony Hotels

* Per person • Double occupancy • Tax not included.



mel r. hertz
MBA, CFP
Certified Financial Planner

Pacific Tower, Suite 2800
1001 Bishop Street
Honolulu, Hawaii 96813
(808) 599-4504

Retirement Plans
•
Investment Management
Consulting
•
Life and Disability
Insurance
•
Charitable Remainder
Trusts



Securities Offered Through IFG Network Securities, Inc. (IFG), Members NASD and SIPC. Financial Planning Services Through Associated Financial Planners, Inc. (AFP), Registered Investment Adviser. mel r hertz is a registered principal of IFG and a registered representative of its affiliate AFP, both of which are otherwise unaffiliated with derand.

A REVIEW OF THE ANOMALOUS ORIGIN OF THE LEFT CORONARY ARTERY FROM THE ANTERIOR SINUS OF VALSALVA: IS PREVENTION POSSIBLE

(Continued from page 297)

right coronary artery. However, all 5 of these cases were associated with sudden death.

A final possible mechanism is coronary vasospasm. Maddoux and associates documented a single case of coronary vasospasm in an anomalous left coronary artery that led to myocardial ischemia²⁴.

It is important to keep in mind that all of these theories could be valid. Sudden death in patients with anomalous left coronary arteries cannot be explained by any one element alone; all of the aforementioned theories might play a role. The final pathway from ischemia to sudden death is commonly ventricular fibrillation²⁵.

It is essential to evaluate any patient with chest pain unattributable to typical causation because of the possible presence of a precarious anomalous left coronary artery. A workup needs to include a base ECG and a radionuclide-scanning exercise stress test to locate regional perfusion deficits. In several cases, this has been helpful in revealing an anomalous left coronary artery²⁶. However, several authors have reported myocardial infarction/sudden death in patients with anomalous left coronary arteries despite previous negative exercise stress tests^{14,27,28}.

Although anomalous left coronary arteries have been identified by echocardiography, angiography eventually will be necessary for a definitive diagnosis^{7,26}. It is essential to define the exact course of the anomalous left coronary artery for both prognostic and surgical reasons. There are 3 components of direction in an anomalous vessel: Sagittal, transverse, and coronal planes. Therefore, 2 views are necessary during angiography; the 2 most informative views are the right anterior oblique and the lateral projections²⁹.

The symptomatic young adult with an anomalous left coronary artery is of primary concern, whereas the same condition in adults is far less likely to result in sudden death. However, it is recommended that all patients undergo surgical correction. A thallium exercise-stress test should be performed. If positive, the anomalous vessel should be bypassed. However, if negative, the patient should be followed over time (Figure 8)²⁶.

The surgical alternatives are: Ostioplasty, separate coronary artery bypass grafting (CABG) to the left anterior descending and circumflex arteries, CABG to the left coronary artery and relocation of the ostium. Enlarging the ostium should be attempted only if the ischemia involved with an anomalous left coronary artery was caused by the slit-like ostium and the acute angle takeoff of the vessel. If the mechanism of ischemia is compression between the great vessels, such an operation would be ineffective. In a review of the literature, there were 3 documented cases of ostioplasty^{13,32,33}. In each case, the patients improved postoperatively. In one of the cases, there was a 7-year follow-up on the patient which corroborated the success of the procedure.

Treatment with bypass grafting has received equal success. Sacks used a single graft to the left coronary artery and said this procedure would cause less restriction of flow than to apply separate grafts to the left anterior descending and circumflex arteries²³. Nevertheless, there also have been many cases of separate bypass grafts that were extremely successful in alleviating symptoms of myocardial ischemia^{7,11,24,26,29,33,34,35,36,37}. These results suggest that compression of

the anomalous left coronary artery plays a significant role in myocardial ischemia.

Mustafa and colleagues suggest an innovative alternative²². The ostium of the left coronary artery is incised and separated from the anterior sinus of Valsalva and the incision extended to above the intercoronary commissure up to the midpoint of the left coronary sinus. At this point, the intima of the left coronary artery and the aorta are joined and the intercoronary commissure reattached to the aorta (Figure 8). That patient did well postoperatively with a normal cardiac catheterization at 1- and 2-year follow-ups²². This procedure also was performed by Donaldson and associates with similar results⁹.

Despite the success of these surgical techniques, it is difficult to judge which procedure is superior. There are no long-term follow-up results. The theoretical efficacy of one operation as compared to another would depend on which theory of the causation of ischemia in anomalous left coronary arteries was correct. The success of various surgical procedures supports the possibility that the ischemia could be caused by a combination of factors.

The ideal remedy would be primary prevention. This concept might appear to be far-fetched; however, as a result of significant advances in embryology and early detection of this anomaly, a preventive modality might loom. The first step toward achieving prevention must lie in understanding how the coronary arteries are formed and what circumstances can result in anomalies. If the causes of coronary anomalies can be discovered and techniques for predevelopmental detection developed, it is quite possible that someday such anomalies might be prevented.

Conclusion

Anomalous origin of the left coronary artery from the anterior sinus of Valsalva can be a serious entity that must be kept in mind in the differential diagnosis of any chest pain otherwise unexplainable. The anomaly is important in terms of the surgical implications, despite its rare occurrence. A high index of suspicion, a correct diagnosis and prompt surgical intervention could prevent sudden death. It is hoped when the embryology of coronary arteries and the causation of anomalies become understood in greater depth, a cost-effective means might be developed for the in-utero detection and prevention of these anomalies.

REFERENCES

- 1) Angelini P. Normal and anomalous coronary arteries: Definitions and classifications. *Tex Heart J*. 1988;117(2):418-434.
- 2) Chaitman BR, Lesperance J, Saltiel J, Bourassa MG. Clinical, angiographic, and hemodynamic findings in patients with anomalous origin of the coronary arteries. *Circulation*. 1975;53:122-135.
- 3) Kimbris D, Iskandrian AS, Segal B, Bemis CE. Anomalous aortic origin of coronary arteries. *Circulation*. 1978;58(4):606-615.
- 4) Alexander RW, Griffith GC. Anomalies of the coronary arteries and their significance. *Am Heart J*. 1956;14: 800- 805.
- 5) Click RL, Holmes DR, Vlietstra RE, Kosinski AS, Dronmal RA. Anomalous coronary arteries: Location, degree of atherosclerosis and effect on survival—a report from the coronary artery surgery study. *JACC*. 1989;13(3):531-537.
- 6) Page HL, Engel HJ, Campbell WB, Thomas CS Jr. Anomalous origin of the left circumflex coronary artery. Recognition, angiographic demonstration and clinical significance. *Circulation*. 1974;50: 768.
- 7) Liberthson RR, Dinsmore RE, Fallon JT. Aberrant coronary artery origin from the aorta. Report of 18 patients, review of the literature and delineation of natu-

- ral history and management. *Circulation*. 1979;59(4):748-754.
- 8) Baltaxe HA, Wixson D. The incidence of congenital anomalies of the coronary arteries in the adult population. *Radiology*. 1974;122: 47.
 - 9) Donaldson RM, Raphael M, Radley-Smith R, Yacoub MH, Ross DN. Angiographic identification of primary coronary anomalies causing impaired myocardial perfusion. *Cathet Cardiovasc Diagn*. 1983;9:237-249.
 - 10) Engel HJ, Torres C, Page HL. Major variations in anatomical origin of the coronary arteries. *Cath Cardiovasc Diagn*. 1975;1:157.
 - 11) Moodie DS, Gill C, Loop FD, Sheldon WC. Anomalous left main coronary artery originating from the right sinus of valsalva. Pathophysiology, angiographic definition and surgical approaches. *J Thorac Cardiovasc Surg*. 1980;80:198-205.
 - 12) Kimbris D. Anomalous origin of the left main coronary artery from the right sinus of valsalva. *Am J Cardiol*. 1985;55: 756-769.
 - 13) Cheitlin MD, DeCastro CM, McAllister HA. Sudden death as a complication of anomalous left coronary origin from the anterior sinus of valsalva. A not-so-minor anomaly. *Circulation*. 1974;50:780-787.
 - 14) Murphy DA, Roy DL, Sohal M, Chandler BM. Anomalous origin of left main coronary artery from anterior sinus of valsalva with myocardial infarction. *J Thorac Cardiovasc Surg*. 1978;75:282-285.
 - 15) Roberts WC, Kragel AH. Anomalous origin of either the right or left main coronary artery from the aorta without coursing of the anomalistically arising artery between aorta and pulmonary trunk. *Am J Cardiol*. 1988;62:1263-1267.
 - 16) Pachinger OM, VandenHoven P, Judkins MP. Single coronary artery—a cause of angina pectoris. *Eur J Cardiol*. 1974;2:161.
 - 17) Roberts WC, Diccio BC, Waller BF, Kishel JC, McManus BM, Dawson SL, Hunsaker JC, Luke JL. Origin of the left main from the right coronary artery of from the right aortic sinus with intramyocardial tunneling to the left side of the heart via the ventricular septum: The case against clinical significance of myocardial bridge or coronary tunnel. *Am Heart J*. 1982;104(2):303-305.
 - 18) Pedal I. Aortale ursprungsanomalie einer koronararterie. *Dtsch Med Wochenschr*. 1976;101:1601.
 - 19) Benson PA, Lack AR. Anomalous aortic origin of the left coronary artery. *Arch Pathol*. 1968;86:214.
 - 20) Corday E. Compression of the left coronary artery during *cor pulmonale*. Proceedings of Second World Congress of Cardiology, Washington DC. 1954;130.
 - 21) Virmani R, Chun PKC, Goldstein RE, Robinowitz M, McAllister HA. Acute takeoffs of the coronary arteries along the aortic wall and congenital coronary ostial valve-like ridges: Association with sudden death. *JACC*. 1984;3(3):766-771.
 - 22) Mustafa I, Gula G, Radley-Smith R, Durrer S, Yacoub M. Anomalous origin of the left coronary artery from the anterior sortic sinus: A potential cause of sudden death. Anatomic characterization and surgical treatment. *J Thorac Cardiovasc Surg*. 1981;82:297-300.
 - 23) Sacks JH, Londe SP, Rosenbluth A, Zalis EG. Left main coronary artery bypass for aberrant (aortic) intramural left coronary artery. *J Thorac Cardiovasc Surg*. 1977;73:733-737.
 - 24) Maddoux GL, Goss JE, Ramo BW, Raff GL, Heuser RR, Shadoff N, Leatherman GF, Blake K, Wilson JN, Deane WM, Hoyt TW, Fowler BN, Gerety RL, Sansonetti D. Angina and vasospasm at rest in a patient with an anomalous left coronary system. *Cath Cardiovasc Diagn*. 1989;16:95-98.
 - 25) Jokl E, McClellan JT, Ross GD. Congenital anomaly of left coronary artery in young athletes. *JAMA*. 1962;182(5):572-573.
 - 26) Cohen AJ, Grishkin BA, Helsel RA, Head HD. Surgical therapy in the management of coronary anomalies: emphasis on utility of internal mammary artery grafts. *Ann Thorac Surg*. 1989;47:630-637.
 - 27) Vander Sande FH, Vandermaast MF, Amsel BJ, Verrelst PA, Walter PJ, Bossaert LL. Cardiac arrest due to aberrant left coronary artery originating from the right sinus of valsalva in a young girl. *Intensive Care Med*. 1989;15:135-136.
 - 28) Barth CW, Roberts WC. Left main coronary artery originating from the right sinus of valsalva and coursing between the aorta pulmonary trunk. *JACC*. 1986;7(2):366-373.
 - 29) Ishikawa T, Brandt PWT. Anomalous origin of the left main coronary artery from the right anterior aortic sinus: angiographic definition of anomalous course. *Am J Cardiol*. 1985;55:770-776.
 - 30) Liberthson RR, Dinsmore RE, Bharati S, Rubenstein JJ, Caulfield J, Wheeler EO, Harthorne JW, Lev M. Aberrant coronary artery origin from the aorta. *Circulation*. 1974;50:774-779.
 - 31) Serota H, Barth CW, Seuc CA, Vandormael M, Aguirre F, Kern MJ. Rapid identification of the course of anomalous coronary arteries in adults: the "dot and eye" method. *Am J Cardiol*. 1990;65:891-898.
 - 32) Grey DP, Koster JK Jr., Farrell PW, Schrank JP. Surgical correction of anomalous left coronary artery from the anterior sinus of valsalva. *Tex Heart Inst J*. 1984;11(2):182-186.
 - 33) Davia JE, Green DC, Cheitlin MD, DeCastro C, Brott WH. Anomalous left coronary artery origin from the right coronary sinus. *Am Heart J*. 1984;108:165-166.
 - 34) Molajo AO, Bray CL, Beton DC, Moussalli H, Prescott MC, Testa HJ. Surgical implications of anomalous aortic origin of the left coronary artery. *J Cardiovascular Surg*. 1989;30:702-705.
 - 35) Bett JHN, O'Brien MF, Murray PJS. Surgery for anomalous origin of the right coronary artery. *Br Heart J*. 1985;53:459-461.
 - 36) Iskandrian AS, Hakki A, Bemis CE. Myocardial ischemia in a patient with anomalous origin of the left main coronary artery. *Cath Cardiovasc Diagn*. 1986;12:48-50.
 - 37) Placentra E, Kornberg B, D'Alonzo A. Anomalous origin of the left coronary artery. *JAMA*. 1989;89(4):506-510.
 - 38) Lauridson JR. Sudden death and anomalous origin of the coronary arteries from the aorta. A case report and review. *Am J Forensic Med Path*. 1988;9(3):236-240.
 - 39) Bogers AJJC, Gittenberger-de Groot AC, Dubbeldam JA, Huysmans HA. The inadequacy of existing theories on development of the proximal coronary arteries and their connexions with the arterial trunks. *Int J Cardiol*. 1988;20:117-123.
 - 40) Bogers AJ, Gittenberger-de Groot AC, Poelmann RE, Peault BM, Huysmans HA. Development of the origin of the coronary arteries, a matter of ingrowth or outgrowth? *Anat Embryol (Berlin)*. 1989;180(5):427-437.
 - 41) Hutchins GM, Kessler-Hanna A, Moore GW. Development of the coronary arteries in the embryonic human heart. *Circ*. 1988;77(6):1250-1257.
 - 42) Kinter AR. Anomalous origin and course of the left coronary artery. *Archives of Pathology*. 1931;586-589.
 - 43) Sanes S. Anomalous origin and course of the left coronary artery in a child. *Am Heart J*. 1978;14: 219-226.
 - 44) Nicod JL. Anomalie coronaire et mort subite. *Cardiologia*. 20: 172. 1952.
 - 45) Dent ED, Fisher RS. Single coronary artery: report of two cases. 1955;1024-1030.
 - 46) Jokl E, McClellan JT, Williams WC, Gouze FJ, Bartholomew RD. Congenital anomaly of left coronary artery in young athletes. *Cardiologia*. 1966;49:253-258.
 - 47) Cohen LS, Shaw LD. Fatal myocardial infarction in an 11-year-old boy associated with a unique coronary artery anomaly. *Am J Cardiol*. 1967;19:420-424.
 - 48) McClellan JT, Jokl E. Congenital anomalies of coronary arteries as cause of sudden death associated with physical exertion. *Am J Clin Path*. 1968;50(2):229-233.
 - 49) Adams RC, Sato K. Sudden death during exercise in an 11- year-old boy. *J Pediatr*. 1974;85(5):737.
 - 50) Schaumburg H, Simonsen J. Sudden death due to congenital malformation of the left coronary artery: a case report. *Forensic Sci Int*. 1978;12: 83-85.
 - 51) Lynch P. Soldiers, sport, and sudden death. *Lancet*. 1980;1:1235-1237.
 - 52) Tsung SH, Huang TY, Chang HH. Sudden death in young athletes. *Arch Pathol Lab Med*. 1982;106: 168-170.
 - 53) Betend B, Billet P, Moreau P, David L. Origine aortique anormale de l'artere coronaire gauche. A propos de la mort subite d'un adolescent. *Arch Fr Pediatr*. 1983;40:479-481.
 - 54) Mitsudo SM, Hirschman AM, Young D. Anomalous origin of the left coronary artery from the right coronary sinus. A cause of sudden death. *NY State J Med*. 1984;402-405.
 - 55) Topaz O, Edwards JE. Pathologic features of sudden death in children, adolescents, and young adults. *Chest*. 1974;87(4):476-482.
 - 56) Daisley H. Sudden death associated with anomalous origin of left coronary artery. *Southern Med J*. 1988; 81(7):944.
 - 57) Mahowald JM, Blieden LC, Coe JI, Edwards JE. Ectopic origin of a coronary artery from the aorta. Sudden death in 3 of 23 patients. *Chest*. 1986;89(5): 668-672.
 - 58) Phillips M, Robinowitz M, Higgins JR, Boran RJ, Reed T, Virmani R. Sudden cardiac death in air force recruits. A 20-year review. *JAMA*. 1986;256(19):2696-2699.
 - 59) Sande JHV, Vandermaast MF, Amsel BJ, Verrelst PA, Walter PJ, Bossaert LL. Cardiac arrest due to aberrant left coronary artery originating from the right sinus of valsalva in a young girl. *Intensive Care Med*. 1989;15:135-136.
 - 60) Kragel AH, Roberts WC. Anomalous origin of either the right or left main coronary artery from the aorta of either the right or left main coronary artery from the aorta with subsequent coursing between aorta and pulmonary trunk: analysis of 32 necropsy cases. *Am J Cardio*. 1988;62:771-777.